



PART-1 (PHYSICS)
JEE Advanced 2017

Q1. A block of mass M has a circular cut with a frictionless surface as shown. The block rests on the horizontal frictionless surface of a fixed table. Initially the right edge of the block is at $x = 0$, in a coordinate system fixed to the table. A point mass m is released from rest at the topmost point of the path as shown and it slides down. When the mass loses contact with the block, its position is x and the velocity is v . At that instant, which of the following options is/are correct?



- (A) The velocity of the point mass m is: $v = \sqrt{\frac{2gR}{1 + \frac{m}{M}}}$
- (B) The velocity of the block M is: $V = -\frac{m}{M} \sqrt{2gR}$
- (C) The position of the point mass is: $x = -\sqrt{2} \frac{mR}{M + m}$
- (D) The x component of displacement of the center of mass of the block M is: $-\frac{mR}{M + m}$


Q2. In the circuit shown, $L = 1\mu\text{H}$, $C = 1\mu\text{F}$ and $R = 1\text{k}\Omega$. They are connected in series with an a.c. source $V = V_0 \sin \omega t$ as shown. Which of the following options is/are correct?



- (A) At $\omega \sim 0$ the current flowing through the circuit becomes nearly zero
- (B) At $\omega \gg 10^6 \text{ rad. s}^{-1}$, the circuit behaves like a capacitor
- (C) The frequency at which the current will be in phase with the voltage is independent of R

(D) The current will be in phase with the voltage if $\omega = 10^4 \text{ rad. s}^{-1}$

Q3. A block M hangs vertically at the bottom end of a uniform rope of constant mass per unit length. The top end of the rope is attached to a fixed rigid support at O. A transverse wave pulse (Pulse 1) of wavelength λ_0 is produced at point O on the rope. The pulse takes time T_{OA} to reach point A. If the wave pulse of wavelength λ_0 is produced at point A (Pulse 2) without disturbing the position of M it takes time T_{AO} to reach point O.



Which of the following options is/are correct?

- (A) The velocities of the two pulses (Pulse 1 and Pulse 2) are the same at the midpoint of rope
- (B) The velocity of any pulse along the rope is independent of its frequency and wavelength
- (C) The wavelength of Pulse 1 becomes longer when it reaches point A
- (D) The time $T_{AO} = T_{OA}$

Q4. A flat plate is moving normal to its plane through a gas under the action of a constant force F. The gas is kept at a very low pressure. The speed of the plate v is much less than the average speed u of the gas molecules. Which of the following options is/are true?

- (A) The pressure difference between the leading and trailing faces of the plate is proportional to uv
- (B) The plate will continue to move with constant non-zero acceleration, at all times
- (C) At a later time external force F balances the resistive force
- (D) The resistive force experienced by the plate is proportional to v

Q5. A human body has a surface area of approximately 1m^2 . The normal body temperature is 10 K above the surrounding room temperature T_0 . Take the room temperature to be $T_0 = 300\text{K}$. For $T_0 = 300\text{K}$, the value of $\sigma T_0^4 = 460\text{Wm}^{-2}$ (where σ is the Stefan-Boltzmann constant). Which of the following options is/are correct?

- (A) Reducing the exposed surface area of the body (e.g. by curling up) allows humans to maintain the same body temperature while reducing the energy lost by radiation
- (B) If the body temperature rises significantly then the peak in the spectrum of electromagnetic radiation emitted by the body would shift to longer wavelengths
- (C) The amount of energy radiated by the body in 1 second is close to 60 Joules

(D) If the surrounding temperature reduces by a small amount $\Delta T_0 \ll T_0$, then to maintain the same body temperature the same (living) human being needs to radiate $\Delta W = 4\sigma T_0^3 \Delta T_0$ more energy per unit time.

Q6. For an isosceles prism of angle A and refractive index μ , it is found that the angle of minimum deviation $\delta m = A$. Which of the following options is/are correct?


(A) At minimum deviation, the incident angle i_1 and the refracting angle r_1 at the first refracting surface are related by $r_1 = (i_1 / 2)$

(B) For this prism, the refractive index μ and the angle of prism A are related as $A = \frac{1}{2} \cos^{-1} \left(\frac{\mu}{2} \right)$

(C) For the angle of incidence $i_1 = A$, the ray inside the prism is parallel to the base of the prism

(D) For this prism, the emergent ray at the second surface will be tangential to the surface when the angle of incidence at the first surface is $i_1 = \sin^{-1} \left[\sin A \sqrt{4 \cos^2 \frac{A}{2} - 1} - \cos A \right]$

Q7. A circular insulated copper wire loop is twisted to form two loops of area A and $2A$ as shown in the figure. At the point of crossing the wires remain electrically insulated from each other. The entire loop lies in the plane (of the paper). A uniform magnetic field \vec{B} points into the plane of the paper. At $t = 0$, the loop starts rotating about the common diameter as axis with a constant angular velocity ω in the magnetic field. Which of the following options is/are correct?



(A) The amplitude of the maximum net emf induced due to both the loops is equal to the amplitude of maximum emf induced in the smaller loop alone


(B) The rate of change of the flux is maximum when the plane of the loops is perpendicular to plane of the paper

(C) The net emf induced due to both the loops is proportional to $\cos \omega t$

(D) The emf induced in the loop is proportional to the sum of the areas of the two loops

Q8. An electron in a hydrogen atom undergoes a transition from an orbit with quantum number n_i to another with quantum number n_f . V_i and V_f are respectively the initial and final potential energies of the electron. If $\frac{V_i}{V_f} = 6.25$ then the smallest possible n_f is _____.

Q9. A monochromatic light is travelling in a medium of refractive index $n = 1.6$. It enters a stack of glass layers from the bottom side at an angle $\theta = 30^\circ$. The interfaces of the glass layers are parallel to each other. The refractive indices of different glass layers are monotonically decreasing as $n_m = n - m\Delta n$, where n_m is the refractive index of the m th slab and $\Delta n = 0.1$ (see the figure). The ray is refracted out parallel to the interface between the $(m-1)^{\text{th}}$ and m^{th} slabs from the right side of the stack. What is the value of m ?



Q10. ^{131}I is an isotope of Iodine that β decays to an isotope of Xenon with a half-life of 8 days. A small amount of a serum labelled with ^{131}I is injected into the blood of a person. The activity of the amount of ^{131}I injected was 2.4×10^5 Becquerel (Bq). It is known that the injected serum will get distributed uniformly in the blood stream in less than half an hour. After 11.5 hours, 2.5 ml of blood is drawn from the person's body, and gives an activity of 115 Bq. The total volume of blood in the person's body, in liters is approximately.

(you may use $e^x \approx 1 + x$ for $|x| \ll 1$ and $\ln 2 \approx 0.7$).

Q11. A stationary source emits sound of frequency $f_0 = 492\text{Hz}$. The sound is reflected by a large car approaching the source with a speed of 2 ms^{-1} . The reflected signal is received by the source and superposed with the original. What will be the beat frequency of the resulting signal in Hz? (Given that the speed of sound in air is 330 ms^{-1} and the car reflects the sound at the frequency it has received).

Q12. A drop of liquid of radius $R = 10^{-2} \text{ m}$ having surface tension $S = \frac{0.1}{4\pi} \text{ Nm}^{-1}$ divides itself into K identical drops. In this process the total change in the surface energy $\Delta U = 10^{-3} \text{ J}$. If $K = 10^\alpha$ then the value of α is _____.



Answer Q.13, Q.14 and Q.15 by appropriately matching the information given in the three columns of the following table.

A charged particle (electron or proton) is introduced at the origin ($x = 0, y = 0, z = 0$) with a given initial velocity \vec{v} . A uniform electric field \vec{E} and a uniform magnetic field \vec{B} exist everywhere. The velocity, \vec{v} electric field and \vec{E} magnetic field \vec{B} are given in columns 1, 2 and 3, respectively. The quantities E_0, B_0 are positive in magnitude.

Column 1		Column 2		Column 3	
(I)	Electron with $\vec{v} = 2 \frac{E_0}{B_0} \hat{x}$	(i)	$\vec{E} = E_0 \hat{z}$	(P)	$\vec{B} = -B_0 \hat{x}$
(II)	Electron with $\vec{v} = \frac{E_0}{B_0} \hat{y}$	(ii)	$\vec{E} = E_0 \hat{y}$	(Q)	$\vec{B} = B_0 \hat{x}$
(III)	Proton with $\vec{v} = 0$	(iii)	$\vec{E} = E_0 \hat{x}$	(R)	$\vec{B} = B_0 \hat{y}$
(IV)	Proton with $\vec{v} = 2 \frac{E_0}{B_0} \hat{x}$	(iv)	$\vec{E} = E_0 \hat{x}$	(S)	$\vec{B} = B_0 \hat{z}$

Q13. In which case will the particle move in a straight line with constant velocity?

- (A) (IV) (i) (S)
- (B) (III) (ii) (R)
- (C) (III) (iii) (P)
- (D) (II) (iii) (S)

Q14. In which case would the particle move in a straight line along the negative direction of y-axis (i.e., move along $-\hat{y}$)?

- (A) (III) (ii) (P)
- (B) (III) (ii) (R)
- (C) (IV) (ii) (S)
- (D) (II) (iii) (Q)

Q15. In which case will the particle describe a helical path with axis along the positive z direction?

- (A) (III) (iii) (P)
- (B) (II) (ii) (R)



(C) (IV) (ii) (R)

(D) (IV) (i) (S)

Answer Q.16, Q.17 and Q.18 by appropriately matching the information given in the three columns of the following table.

An ideal gas is undergoing a cyclic thermodynamic process in different ways as shown in the corresponding P-V diagrams in column 3 of the table. Consider only the path from state 1 to state 2. W denotes the corresponding work done on the system. The equations and plots in the table have standard notations as used in thermodynamic processes. Here γ is the ratio of heat capacities at constant pressure and constant volume. The number of moles in the gas is n .

Column 1		Column 2		Column 3	
(I)	$W_{1 \rightarrow 2} = \frac{1}{\gamma-1}(P_2V_2 - P_1V_1)$	(i)	Isothermal	(P)	
(II)	$W_{1 \rightarrow 2} = -PV_2 + PV_1$	(ii)	Isochoric	(Q)	
(III)	$W_{1 \rightarrow 2} = 0$	(iii)	Isobaric	(R)	
(IV)	$W_{1 \rightarrow 2} = -nRT \ln\left(\frac{V_2}{V_1}\right)$	(iv)	Adiabatic	(S)	

Q16. Which of the following options is the only correct representation of a process in which?

$$\Delta U = \Delta Q - P\Delta V ?$$

(A) (II) (iv) (R)

(B) (III) (iii) (P)

(C) (II) (iii) (P)

(D) (II) (iii) (S)

Q17. Which one of the following options correctly represents a thermodynamic process that is used as a correction in the determination of the speed of sound in an ideal gas?



(A) (I) (iv) (Q)

(B) (III) (iv) (R)

(C) (I) (ii) (Q)

(D) (IV) (ii) (R)

18. Which one of the following options is the correct combination?

(A) (IV) (ii) (S)

(B) (III) (ii) (S)

(C) (II) (iv) (R)

(D) (II) (iv) (P)

Copyright ©Jagranjosh.com

All rights reserved. No part or the whole of this eBook may be copied, reproduced, stored in retrieval system or transmitted and/or cited anywhere in any form or by any means (electronic, mechanical, photocopying, recording or otherwise), without the written permission of the copyright owner. If any misconduct comes in knowledge or brought in notice, strict action will be taken.